


Claims

1. An immunoeffector compound having the following structure:



wherein, X is selected from the group consisting of O and S at the axial or equatorial position; Y is selected from the group consisting of O and NH; n, m, p and q are integers from 0 to 6; R₁, R₂ and R₃ are the same or different and are normal fatty acyl residues having from 1 to about 20 carbon atoms and where one of R₁, R₂ or R₃ is optionally hydrogen; R₄ and R₅ are the same or different and are selected from the group consisting of H and methyl; R₆ and R₇ are the same or different and are selected from the group consisting of H, hydroxy, alkoxy, phosphono, phosphonooxy, sulfo, sulfooxy, amino, mercapto, cyano, nitro, formyl and carboxy, and esters and amides thereof; and R₈ and R₉ are the same or different and are selected from the group consisting of phosphono and H, and at least one of R₈ and R₉ is phosphono.

2. The compound of claim 1, wherein R₆ is carboxy.

3. The compound of claim 2, wherein X is O; Y is O; n, m, p and q are 0; R₁, R₂ and R₃ are normal fatty acyl residues having 10 carbon atoms; R₄, R₅ and R₇ are H; R₈ is phosphono; R₉ is H; R₁, R₂ and R₃ are each attached to a stereogenic center having an *R* configuration; and R₅ is attached to a stereogenic center having an *S* configuration.

4. The compound of claim 2, wherein X is O; Y is O; n, m, p and q are 0; R₁, R₂ and R₃ are normal fatty acyl residues having 12 carbon atoms; R₄, R₅ and R₇ are H; R₈ is phosphono; R₉ is H; R₁, R₂ and R₃ are each attached to a stereogenic center having an *R* configuration; and R₅ is attached to a stereogenic center having an *S* configuration.

5. The compound of claim 2, wherein X is O; Y is O; n, m, p and q are 0; R₁, R₂ and R₃ are normal fatty acyl residues having 10 carbon atoms; R₄, R₅ and R₇ are H; R₈ is phosphono; R₉ is H; R₁, R₂ and R₃ are each attached to a stereogenic center having an *R* configuration; and R₅ is attached to a stereogenic center having an *R* configuration.

6. The compound of claim 2, wherein X is O; Y is O; n, m, p and q are 0; R₁, R₂ and R₃ are normal fatty acyl residues having 8 carbon atoms; R₄, R₅ and R₇ are H; R₈ is phosphono; R₉ is H; R₁, R₂ and R₃ are each attached to a stereogenic center having an *R* configuration; and R₅ is attached to a stereogenic center having an *S* configuration.

7. The compound of claim 1, wherein R₆ is H.

8. The compound of claim 7, wherein X is O; Y is O; n is 2; m, p and q are 0; R₁, R₂ and R₃ are normal fatty acyl residues having 14 carbon atoms; R₄, R₅ and R₇ are H; R₈ is phosphono; R₉ is H; and R₁, R₂ and R₃ are each attached to a stereogenic center having an *R* configuration.

9. The compound of claim 7, wherein X is O; Y is O; n is 1, m and p are 0; q is 1; R₁, R₂ and R₃ are normal fatty acyl residues having 10 carbon atoms; R₄ and R₅ are

H; R₇ is carboxy; R₈ is phosphono; R₉ is H; and R₁, R₂ and R₃ are each attached to a stereogenic center having an *R* configuration.

10. The compound of claim 7, wherein X is O; Y is O; m, n, p and q are 0; R₁, R₂ and R₃ are normal fatty acyl residues having 14 carbon atoms; R₄, R₅ and R₇ are H; R₈ is phosphono; R₉ is H; and R₁, R₂ and R₃ are each attached to a stereogenic center having an *R* configuration.

11. The compound of claim 7, wherein X is O; Y is O; m, n, p and q are 0; R₁, R₂ and R₃ are normal fatty acyl residues having 10 carbon atoms; R₄, R₅ and R₇ are H; R₈ is phosphono; R₉ is H; and R₁, R₂ and R₃ are each attached to a stereogenic center having an *R* configuration.

12. The compound of claim 7, wherein X is O; Y is O; m, p and q are 0; n is 1; R₁, R₂ and R₃ are normal fatty acyl residues having 14 carbons; R₄, R₅ and R₇ are H; R₈ is phosphono; R₉ is H; and R₁, R₂ and R₃ are each attached to a stereogenic center having an *R* configuration.

13. The compound of claim 1, wherein R₆ is hydroxy.

14. The compound of claim 13, wherein X is O; Y is O; m, n and q are 0; p is 1; R₁, R₂ and R₃ are normal fatty acyl residues having 12 carbon atoms; R₄ and R₅ are H; R₇ is H; R₈ is phosphono; and R₉ is H; R₁, R₂ and R₃ are each attached to a stereogenic center having an *R* configuration; and R₅ is attached to a stereogenic center having an *S* configuration.

15. The compound of claim 13, wherein X is O; Y is O; m and q are 0; n and p are 1; R₁, R₂ and R₃ are normal fatty acyl residues having 10 carbon atoms; R₄, R₅ and R₇ are H; R₈ is phosphono; R₉ is H; R₁, R₂ and R₃ are each attached to a stereogenic center

having an *R* configuration; and R_5 is attached to a stereogenic center having an *S* configuration.

16. The compound of claim 13, wherein X is O ; Y is O ; m , n and q are 0 ; p is 2 ; R_1 , R_2 and R_3 are normal fatty acyl residues having 10 carbon atoms; R_4 , R_5 and R_7 are H ; R_8 is phosphono; R_9 is H ; R_1 , R_2 and R_3 are each attached to a stereogenic center having an *R* configuration; and R_5 is attached to a stereogenic center having an *S* configuration.

17. The compound of claim 13, wherein X is O ; Y is O ; m , n and q are 0 ; p is 1 ; R_1 , R_2 and R_3 are normal fatty acyl residues having 14 carbon atoms; R_4 , R_5 and R_7 are H ; R_8 is phosphono; R_9 is H ; R_1 , R_2 and R_3 are each attached to a stereogenic center having an *R* configuration; and R_5 is attached to a stereogenic center having an *R* configuration.

18. The compound of claim 13, wherein X is O ; Y is O ; m , n and q are 0 ; p is 1 ; R_1 , R_2 and R_3 are normal fatty acyl residues having 14 carbon atoms; R_4 , R_5 and R_7 are H ; R_8 is phosphono; R_9 is H ; R_1 , R_2 and R_3 are each attached to a stereogenic center having an *R* configuration; and R_5 is attached to a stereogenic center having an *S* configuration.

19. The compound of claim 13, wherein X is O ; Y is O ; m , n and q are 0 ; p is 1 ; R_1 , R_2 and R_3 are normal fatty acyl residues having 11 carbon atoms; R_4 , R_5 and R_7 are H ; R_8 is phosphono; R_9 is H ; R_1 , R_2 and R_3 are each attached to a stereogenic center having an *R* configuration; and R_5 is attached to a stereogenic center having an *S* configuration.

20. The compound of claim 13, wherein X is O ; Y is O ; m , n and q are 0 ; p is 1 ; R_1 , R_2 and R_3 are normal fatty acyl residues having 10 carbon atoms; R_4 , R_5 and R_7 are H ; R_8 is phosphono; R_9 is H ; R_1 , R_2 and R_3 are each attached to a stereogenic center having an *R* configuration; and R_5 is attached to a stereogenic center having an *S* configuration.

21. The compound of claim 1, wherein X is O ; Y is O ; m , n , p and q are 0 ; R_1 , R_2 and R_3 are normal fatty acyl residues having 10 carbon atoms; R_4 and R_5 are H ; R_6 is

amino carbonyl; R_7 is H; R_8 is phosphono; and R_9 is H; R_1 , R_2 and R_3 are each attached to a stereogenic center having an *R* configuration; and R_5 is attached to a stereogenic center having an *S* configuration.

22. The compound of claim 1, wherein R_1 is hydrogen.
23. The compound of claim 1, wherein R_2 is hydrogen.
24. The compound of claim 1, wherein R_3 is hydrogen.
25. A method for enhancing the immune response of a mammal comprising administering to the mammal an effective amount of a compound of claim 1.
26. A vaccine composition comprising a compound of claim 1, an antigen and a suitable carrier.
27. A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.
28. The composition of claim 27, wherein said pharmaceutically acceptable carrier is an aqueous composition comprising water and one or more surfactants selected from the group consisting of glycodeoxycholate, deoxycholate, sphingomyelin, sphingosine, phosphatidylcholine, 1,2-Dimyristoyl-sn-glycero-3-phosphoethanolamine, L- α -Phosphatidylethanolamine, and 1,2-Dipalmitoyl-sn-glycero-3-phosphocholine, or a mixture thereof.
29. The composition of claim 28, wherein said one or more surfactant is 1,2-Dipalmitoyl-sn-glycero-3-phosphocholine.

30. The composition of claim 28, wherein the molar ratio of said compound to surfactant is from about 10:1 to about 1:25.

Sub. to add
should be
31. The composition of claim 28, wherein the molar ratio of said compound to surfactant is from about 4:1 to about 1:9.

32. The composition of claim 27, wherein said carrier is a stable emulsion comprising a metabolizable oil, one or more surfactants, an antioxidant and a component to make the emulsion isotonic.

33. The composition of claim 32, wherein said stable emulsion comprises 1-10% v/v squalene, 0.9% w/v PLURONIC-F68 block co-polymer, 1.9% w/v egg phosphatidyl choline, 1.75% v/v glycerol and 0.05% w/v α tocopherol.

34. The composition of claim 27 wherein said carrier is a suspension comprising aluminum hydroxide, calcium hydroxide, calcium phosphate or tyrosine adsorbate.

35. The composition of claim 27 wherein said carrier is an aqueous solution or aqueous micellar dispersion comprising triethylamine or triethanolamine.

36. The composition of claim 27 wherein said carrier comprises microspheres or microparticles, and the compound of claim 1 is within the matrix of the microspheres or microparticles or adsorbed thereon.